

## DESCRIPTION

Advanced Protocol® Verified 75 IF is a Constant Nutrition® formulation providing 20% protein for mice and rats in protocols where low dietary estrogenic activity needs to be assured. This diet is formulated using the unique and innovative concept of Constant Nutrition®, paired with the selection of highest quality ingredients to assure minimal inherent biological variation in long-term studies.

### Features and Benefits

- Constant Nutrition® formula helps minimize nutritional variables
- Formulated with 20% protein
- Verified to contain less than 75 ppm total isoflavones (genistein, daidzein and glycitein)
- Provides proper nutrients without affecting outcome in estrogen-sensitive protocols
- Precision processing assures Constant Nutrition® quality

### Product Forms Available

- Oval pellets, 10 mm x 16 mm x 25 mm length (3/8"x5/8"x1")
- Meal (ground pellets), special order

### Other Versions Available

- 5V12: Advanced Protocol Extruded Verified 75 IF Diet

## GUARANTEED ANALYSIS

Crude protein not less than	.20.0%
Crude fat not less than	.4.5%
Crude fiber not more than	.4.0%

## INGREDIENTS

Ground wheat, wheat middlings, corn gluten meal, ground corn, wheat germ, cane molasses, dried beet pulp, soybean oil, calcium carbonate, dicalcium phosphate, monocalcium phosphate, L-lysine, dehulled soybean meal, salt, DL-methionine, dried whey, choline chloride, menadione dimethylpyrimidinol bisulfite, L-tryptophan, pyridoxine hydrochloride, chromium potassium sulfate, potassium chloride, tocopherols, cholecalciferol, vitamin A acetate, magnesium oxide, biotin, dl-alpha tocopheryl acetate, folic acid, thiamin mononitrate, vitamin B<sub>12</sub> supplement, calcium pantothenate, nicotinic acid, riboflavin, zinc oxide, manganese oxide, ferrous carbonate, copper sulfate, zinc sulfate, calcium iodate, cobalt carbonate, sodium selenite.

## FEEDING DIRECTIONS

Feed ad libitum. Plenty of fresh, clean water should be available to the animals at all times.

**Rats-** All rats will eat varying amounts of feed depending on their genetic origin. Larger strains will eat up to 30 grams per day. Smaller strains will eat up to 15 grams per day. Feeders in rat cages should be designed to hold two to three days supply of feed at one time.

**Mice-** Adult mice will eat up to 5 grams of pelleted ration daily. Some of the larger strains may eat as much as 8 grams per day per animal. Feed should be available on a free choice basis in wire feeders above the floor of the cage.

**Hamsters-** Adults will eat up to 14 grams per day.

**LabDiet** ADVANCED PROTOCOL™

## CHEMICAL COMPOSITION<sup>1</sup>

Nutrients <sup>2</sup>		
<b>Protein, %</b>	<b>.20.0</b>	Sulfur, % . . . . .0.24
Arginine, % . . . . .	.0.79	Sodium, % . . . . .0.25
Cystine, % . . . . .	.0.31	Chlorine, % . . . . .0.58
Glycine, % . . . . .	.0.60	Fluorine, ppm . . . . .15
Histidine, % . . . . .	.0.40	Iron, ppm . . . . .220
Isoleucine, % . . . . .	.0.83	Zinc, ppm . . . . .130
Leucine, % . . . . .	.2.17	Manganese, ppm . . . . .130
Lysine, % . . . . .	.1.00	Copper, ppm . . . . .17
Methionine, % . . . . .	.0.66	Cobalt, ppm . . . . .0.63
Phenylalanine, % . . . . .	.0.99	Iodine, ppm . . . . .1.5
Tyrosine, % . . . . .	.0.71	Chromium, ppm . . . . .1.3
Threonine, % . . . . .	.0.62	Selenium, ppm . . . . .0.34
Tryptophan, % . . . . .	.0.24	
Valine, % . . . . .	.0.86	<b>Vitamins</b>
Serine, % . . . . .	.0.93	Carotene, ppm . . . . .2.2
Aspartic Acid, % . . . . .	.1.19	Vitamin K (as menadione), ppm . . . . .4.0
Glutamic Acid, % . . . . .	.4.85	Thiamin Hydrochloride, ppm . . . . .12
Alanine, % . . . . .	.1.23	Riboflavin, ppm . . . . .7.0
Proline, % . . . . .	.1.92	Niacin, ppm . . . . .94
Taurine, % . . . . .	.0.00	Pantothenic Acid, ppm . . . . .18
<b>Fat (ether extract), %</b>	<b>.5.0</b>	Choline Chloride, ppm . . . . .2000
<b>Fat (acid hydrolysis), %</b>	<b>.5.5</b>	Folic Acid, ppm . . . . .2.5
Cholesterol, ppm . . . . .	.0.00	Pyridoxine, ppm . . . . .10
Linoleic Acid, % . . . . .	.2.70	Biotin, ppm . . . . .0.30
Linolenic Acid, % . . . . .	.0.28	B <sub>12</sub> , mcg/kg . . . . .51
Arachidonic Acid, % . . . . .	.0.00	Vitamin A, IU/gm . . . . .15
Omega-3 Fatty Acids, % . . . . .	.0.30	Vitamin D <sub>3</sub> (added), IU/gm . . . . .2.2
Total Saturated Fatty Acids, % . . . . .	.0.86	Vitamin E, IU/kg . . . . .70
Total Monounsaturated		Ascorbic Acid, mg/gm . . . . .—
Fatty Acids, % . . . . .	.1.08	
<b>Fiber (Crude), %</b>	<b>.3.8</b>	<b>Calories provided by:</b>
Neutral Detergent Fiber <sup>3</sup> , % . . . . .	.17.0	Protein, % . . . . .23.035
Acid Detergent Fiber <sup>4</sup> , % . . . . .	.5.5	Fat (ether extract), % . . . . .12.886
<b>Nitrogen-Free Extract</b>		Carbohydrates, % . . . . .64.078
<b>(by difference), %</b>	<b>.55.6</b>	*Product Code
Starch, % . . . . .	.32.9	1. Formulation based on calculated values from the latest ingredient analysis information. Since nutrient composition of natural ingredients varies and some nutrient loss will occur due to manufacturing processes, analysis will differ accordingly.
Glucose, % . . . . .	.0.11	2. Nutrients expressed as percent of ration except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.
Fructose, % . . . . .	.0.10	3. NDF = approximately cellulose, hemicellulose and lignin.
Sucrose, % . . . . .	.2.48	4. ADF = approximately cellulose and lignin.
Lactose, % . . . . .	.0.15	5. Physiological Fuel Value (kcal/gm) = Sum of decimal fractions of protein, fat and carbohydrate (use Nitrogen Free Extract) x 4,9,4 kcal/gm respectively.
<b>Total Digestible Nutrients, %</b>	<b>.77.4</b>	
<b>Gross Energy, kcal/gm</b>	<b>.4.12</b>	
<b>Physiological Fuel Value<sup>5</sup>, kcal/gm</b>	<b>.3.47</b>	
<b>Metabolizable Energy, kcal/gm</b>	<b>.3.23</b>	
<b>Minerals</b>		
<b>Ash, %</b>	<b>.5.4</b>	
Calcium, % . . . . .	.0.87	
Phosphorus, % . . . . .	.0.60	
Phosphorus (non-phytate), % . . . . .	.0.32	
Potassium, % . . . . .	.0.64	
Magnesium, % . . . . .	.0.20	