PMI® Rodent Liquid Diet LD I 02*

Technical Data

DESCRIPTION

Diet LD 102 is a dry powder used to prepare a liquid diet formulated for rodents. It is nutritionally balanced with excellent palatability. This diet is especially designed for use as the control diet when LD 102A is being used as the test diet. LD 102 can be used in other applications where solid diets are not appropriate.

Features and Benefits

- · Nutritionally-balanced
- · Volatile ingredients can be included
- · Easily prepared
- Provides stable nutrients
- Shipped in dry form to simplify storage, shipping and stability
- · Minimal foaming
- Fully suspended
- · Stabilized against microbial growth

Product Forms Available	Catalog #
• Dry Powder	0007553 (530A)
Dry Powder, Irradiated	1815250 (530A)
GUARANTEED ANALYSIS	,
Crude protein not less than	

Crude protein not less than	. 16.00%
Crude fat not less than	. 15.00%
Crude fiber not more than	. 10.00%
Ash not more than	5.00%

*Diet Preparation Instructions: To 771 gms. of water, add 229 gms. Rodent Liquid Diet mix (LD 102). Blend vigorously for 15-30 seconds with a mechanical blender until completely suspended. For best results add water to blender before dry mix.

Additional Considerations:

- For best results a mechanical blender should be used for diet preparation; hand blending does not suspend the diet adequately to avoid some settling out of undissolved ingredients.
- Do not over-blend; excessive mechanical blending creates foaming.

INGREDIENTS

Vitamin-free casein, olive oil, maltodextrin, dextrose, soy fiber, corn oil, suspension colloid, safflower oil, L-cystine, DL-methionine, vitamin A acetate, cholecalciferol, dl-alpha tocopheryl acetate, menadione dimethylpyrimidinol bisulfite (source of vitamin K), ascorbic acid, cyanocobalamin, thiamin mononitrate, riboflavin, calcium pantothenate, nicotinic acid, choline chloride, pyridoxine hydrochloride, folic acid, inositol, p-aminobenzoic acid, biotin, calcium acetate, calcium phosphate, potassium phosphate, sodium phosphate, magnesium sulfate, sodium chloride, manganese sulfate, ferrous fumarate, zinc chloride, cupric sulfate, chromium chloride, sodium fluoride, ammonium molybdate, calcium iodate, sodium selenite.

FEEDING DIRECTIONS

Diet consumption will vary according to animal size and sex. An average rat should consume about 74-109 grams of liquid diet (17-25 grams dry diet) daily. The growth rate of rats maintained on this diet should be similar to that attained by young rats (55-100 grams) maintained on a good quality, nonpurified rodent diet. Mice should consume at least 20 grams of liquid diet per day. Allow new animals an adequate period of time to adjust to their surroundings. After they have adjusted, introduce the liquid diet gradually by offering some of the liquid diet while the regular diet is still present. Gradually decrease the amount of regular diet offered while increasing the amount of liquid diet over a 3-5 day period. Additional time for adjustment may be necessary for the ethanol diets. Prepare the diet daily for best results as bacterial growth will be excessive in 48 hours. Additional water may be provided in separate drinking tubes, but may not be consumed.

For information regarding shelf life please visit www.labdiet.com.

CHEMICAL COMPOSITION'

Nutrients ²	Recon- stituded ²	
Protein, %	4.06	17.7
Arginine, %		0.70
Cystine, %	0.07	0.30
Glycine, %	0.09	0.39
Histidine, %	0.12	0.53
Isoleucine, %	0.22	0.94
Leucine, %	0.39	1.70
Lysine, %	0.33	1.42
Methionine, %	0.15	0.63
Phenylalanine, %	0.22	0.94
Tyrosine, %	0.23	0.99
Threonine, %	0.18	0.77
Tryptophan, %	0.05	0.21
Valine, %	0.26	1.12
Aspartic Acid, %	0.30	1.28
Glutamic Acid, %	0.92	3.98
Fat (ether extract), %	3.9	16.9
Fiber (Crude), %	0.69	3.0
A4: 1		

Minerals	
Calcium, %0.14	0.61
Phosphorus, %0.11	0.47
Potassium, %0.12	0.53
Magnesium, %0.03	0.11
Sulfur, %0.06	0.26
Sodium, %0.06	0.26
Chloride, %0.07	0.30
Fluorine, ppm0.25	1.10
Iron, ppm18	80
Zinc, ppm9.2	40
Manganese, ppm14	59
Copper, ppm2.5	11
Chromium (added), ppm0.60	2.6
Iodine, ppm0.05	0.21
Molybdenum, ppm0.11	0.50
Selenium, ppm0.02	0.07

Vitamins

Vicuilliis	
Vitamin K, ppm0.25	1.1
Thiamin, ppm1.5	6.5
Riboflavin, ppm1.7	7.2
Niacin, ppm7.6	33
Pantothenic Acid, ppm3.9	17
Choline Chloride, ppm265	1150
Folic Acid, ppm0.53	2.3
Pyridoxine, ppm1.4	6.3
Biotin, ppm0.05	0.22
Inositol, ppm25	110
p-aminobenzoic acid, ppm12.5	54
B ₁₂ , mcg/kg25	110
Vitamin A, IU/gm3.0	13
Vitamin D ₃ (added), IU/gm0.40	1.7
Vitamin E, IU/kg30	130
Ascorbic Acid, mg/gm9.0	39

Calories provided by:

Protein, kcal/kg173
Fat, Kcal/kg350
Carbohydrates, kcal/kg477
*Energy Levels used (kcal/gm)
Protein = 4.25; Fat = 9.00; Maltodextrin =
4.00; Ethanol = 7.07 . The protein value is
different than the 4 kcal/gm for protein, as
generally used.

- * 1 kilogram of diet in liquid form, when prepared according to directions, provides 1000 kilocalories (1 kcal per gram).
- * Lieber, CS & LM DeCarli (1982) Alcoholism: Clinical and Experimental Research 6: 523-531. Miller, SS, ME Goldman, CK Erickson & RL Shorey (1980) Psychopharmacology 68: 55-59.
- *Product Code 1.Based on the latest ingredient analysis information.
- 2. Values are based upon the liquid form of the diet when prepared according to directions (230 gm dry powder with 770 gm water).

